



AP 10-0-10 6.8% S



GUARANTEED ANALYSIS

Total Nitrogen (N).....	10.00%
3.0% Water Soluble Organic Nitrogen	
7.0% Urea Nitrogen	
Soluble Potash (K ₂ O).....	10.00%
Sulfur (S).....	6.80%
Derived from: Polyethylene Urea Solution, Urea and Potassium Thiosulfate	

GENERAL INFORMATION

Harvey's 10-0-10 6.8% S is a proprietary liquid fertilizer solution, deriving its slow-release properties from methylene ureas (estimated release pattern is 4 – 8 weeks). **Harvey's 10-0-10 6.8% S** is recommended for Row Crops, Ornamentals and turf, Vegetable Crops, Fruit and Nut Crops. **Harvey's 10-0-10 6.8% S** which is ideal for ground applications as well as drip irrigation systems. It is compatible with other NPK fertilizers, micronutrients and can be tank mixed with most herbicides, insecticides, and fungicides.

PRECAUTION: It is best to apply **Harvey's 10-0-10 6.8% S** in the early morning or late evening to maximize uptake and efficiency and to avoid crop injury. As with the use of any agrichemical, crop injury is always possible. Foliar fertilization is intended as a supplement to a regular fertilization program and may not, by itself, provide all the nutrients normally required by crops or other intended plants. Avoid strong acids or contact with aluminum, mild steel and brass. Jar tests should be performed to confirm compatibility before mixing with other components.

RECOMMENDED RATES: Rates and timing of applications are dependent on local conditions, and should always be made as a result of soil or plant tissue analysis. Best results are achieved when **Harvey's 10-0-10 6.8% S** is used to supplement a fertility program based on soil tests. Ground or aerial systems may be utilized to deliver **Harvey's 10-0-10 6.8% S**. Use sufficient water to ensure thorough coverage. Product may be applied either diluted or undiluted. Higher rates may be applied if done so first on a small test area to determine acceptability.

Soybeans: Apply 1-4 Qts. per acre at early pod formation or Apply 1-4 Qts. per acre at early full flower and repeat in 10 – 14 days.

Corn (Sweet): Apply 1-3 gals. per acre foliarly at V-6 to V-8 stage as a partial side-dress Nitrogen (N) replacement. Apply 1 – 3 gals. per acre foliarly at pre-tassel as a Nitrogen (N) supplement. Apply 1 – 2 gals. per acre foliarly at V1 to R-2 stage with fungicides or insecticides. These applications can be combined with glyphosate.

Cotton: Seedling Stage: Apply 2-4 qts. per acre when first true leaves appear. Alter Seedling Stage: Apply 2-4 Qts. per acre. Boll Development: 2-4 Qts. per acre after bloom, repeat in 7 day intervals.

Tobacco: Apply 1-4 qts. per acre at plant bed stage to near maturity as needed to maintain growth and quality

Grain Sorghum: Apply 1-3gal. per acre after pollination

Cucumbers: Apply 2-4 gals. per acre at early fruit set. Or apply 2-4gals. per acre at early flowering and repeat at 10 – 14 day intervals

Watermelons: Apply 1-2 gals. per acre at early flowering and repeat 7 – 10 days later

Turf, Bent Grass: Apply 4.5 – 6.8 lbs. of Nitrogen per 1,000 sq. ft. per year.

Bermuda Grass (Turf): Apply 9.0 – 11.5 lbs. of Nitrogen per 1,000 sq. ft. per year. May apply in multiple applications of 0.25 lbs. to 0.5 lbs. of Nitrogen

St. Augustine (includes other Warm Season Grass): Apply 2.5 – 3.5 lbs. of Nitrogen per 1,000 sq. ft. per year. May apply in multiple applications as needed.

For all Turf recommendations above, in areas with prolonged growing seasons, areas of high foot traffic or in "grow-in" situations, higher rates may be used.

TANK MIXING AND BLENDING

Dilute with water and blend with other nutrients and pesticides only at the time of application and in the amounts required. Jar tests should be performed to confirm compatibility before mixing with other components. Fill the clean spray or mix tank 3/4 with water, begin agitation, add other materials in the following sequence (unless otherwise directed by their labeling):

1. Add 3/4 total water to the tank.
2. Turn on the agitator or recirculation line.
3. Add prescribed amount of **Harvey's 10-0-10 6.8% S** to the tank.
4. Add any soluble powder.
5. Add liquid flowable materials.
6. Bring water to volume and recirculate before spraying.

CAUTION: Harmful if swallowed. Do not take internally. Avoid contact with eyes, skin, or clothing. Wear appropriate personal protective equipment (goggles and face shield) when handling this product. Do not apply this product in such a manner as to directly expose workers or other persons. If product is being mixed with pesticides, spray adjuvants and/or compatibility agents, follow the "Precautionary Statement/FIRST AID" on the product's labeling

FIRST AID IF SWALLOWED:

- Have person drink large volumes of water (2 to 4 glasses of water).
- Call a poison control center or physician immediately for treatment advice.
- Do not induce vomiting unless told to do so by a poison control center or physician.
- Do not give anything by mouth to an unconscious person.

IF IN EYES:

- Hold eye open and rinse slowly and gently with water for 15 to 20 minutes.
- Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye.
- Call a poison control center or physician for treatment advice.

IF ON SKIN OR CLOTHING:

- Take off contaminated clothing.
- Rinse skin immediately with plenty of water for 15 to 20 minutes.
- Call a poison control center or physician for treatment advice if irritation develops.

WARRANTY—CONDITIONS OF SALE

Our Directions for use of this product are based upon rates believed reliable. Follow directions carefully. Timing and method of application, weather and crop conditions, mixtures with other chemicals not specifically recommended and other influencing factors in the use of this product are beyond the control of the Seller. To the extent consistent with applicable law, Buyer assumes all risks of use, storage and handling of this material not in strict accordance with directions given herewith.

To the extent consistent with applicable law, in no case shall the Manufacturer or the Seller be liable for consequential, special or indirect damages resulting from the use or handling of this product when such use and/or handling is not in strict accordance with directions given herewith. The foregoing is a condition of sale by the Seller and is accepted as such by the Buyer.

STORAGE:

Store at temperatures between 35°F – 90°F. Keep out of reach of children and animals in a cool, dry chemical storage area. Keep container tightly closed. Do not allow water to be introduced into the container contents.

DISPOSAL:

Dispose of empty containers by triple rinsing with detergent solution or puncture and discard empty containers in a landfill in accordance with current Local, State and Federal regulations. Do not contaminate water sources by cleaning of equipment or disposal of waste.

Net Weight:

- Jug: 26lb (11.82kg)
- Cx: 2x2 1/2: 52lb (23.64kg)

Net Contents:

- Jug: 2 1/2 gal (9.46 liters)
- Cx: 2x2 1/2: 5 gal (18.92 liters)

